Parasites in Domestic Cats

Identification, Treatment, Prevention

Cat Man (Booklet's)

Parasites in Domestic Cats
Identification, Treatment, Prevention



Hrsg: Cat Man (Booklet's)
Godshorn (Germany) 2025

Introduction
Parasites as a Health Risk for Domestic Cats

Parasites represent one of the most common health risks for domestic cats and can significantly affect their well-being. Addressing this topic is essential because a parasitic infestation not only causes discomfort and health issues in cats but, in some cases, also poses risks to humans. The range of problems caused by parasites extends from mild irritation to potentially life-threatening diseases. Early detection and treatment are therefore crucial for recovery and long-term health. Regular veterinary check-ups and parasite screening are indispensable for identifying infestations in time and taking appropriate action.

In general, parasites that affect cats are divided into two main categories: **internal** and **external** parasites. Internal parasites live inside the cat's body, such as in the intestines or heart, while external parasites live on the body's surface, like fleas or ticks. Understanding this distinction is essential to recognize the different types of infections and to apply the right treatment and prevention methods.

Internal Parasites – A Detailed Overview

Internal parasites are a common problem in cats and can cause a wide range of often nonspecific symptoms. Studies show that prevalence rates in some feline populations can reach up to 45 percent. These parasites can be either worm-like (helminths) or single-celled organisms (protozoa). The high occurrence demonstrates that most cats will come into contact with internal parasites at some point in their lives.

Internal parasites can interfere with the absorption of nutrients, damage tissues, and weaken the immune system. Kittens, elderly cats, and those with compromised health are particularly at risk. The symptoms can range from mild digestive issues to serious systemic illness, depending on the type and number of parasites present. Regular deworming and preventive care are therefore essential to protect cats from the harmful effects of internal parasites.

Roundworms (Ascarids)

Roundworms are the most common internal parasites in cats, affecting an estimated 25 to 75 percent of all cats, with infection rates even higher among kittens. The two primary species that infect cats are *Toxascaris leonina* and *Toxocara cati*. Adult worms can reach lengths of three to five inches, are cream-colored, and resemble spaghetti in appearance.

Cats can become infected in several ways — by ingesting eggs from contaminated soil or water, by licking contaminated fur or paws, or by eating infected rodents that serve as paratenic hosts. A key route of transmission for *Toxocara cati* is through the mother's milk to her kittens.

Symptoms of roundworm infection vary widely. Adult cats may show no or only mild signs, while kittens often develop diarrhea, vomiting (sometimes containing worms), a swollen belly, dull coat, weight loss, and poor growth. When larvae migrate through the lungs, coughing or breathing difficulties may occur. In severe infestations, anemia can develop, recognizable by pale gums.

Roundworms deprive the cat of nutrients, leading to malnutrition and digestive problems. In kittens, migrating larvae can cause respiratory disease such as pneumonia. Untreated heavy infestations can result in life-threatening anemia or, in extreme cases, intestinal rupture.

Roundworm infection also poses a **zoonotic risk** — meaning it can affect humans. Infection usually occurs through accidental ingestion of eggs from contaminated soil, particularly among children. The larvae can migrate through the human body to organs such as the liver, lungs, brain, and eyes, leading to tissue damage (visceral or ocular larva migrans). In severe cases, this may cause organ impairment or blindness. Early and consistent deworming, combined with good hygiene, is the best way to prevent transmission.

Hookworms (Ancylostomatidae)

Hookworms are also common in dogs but can infect cats as well. These slender, threadlike worms measure less than half an inch in length and attach to the intestinal lining, feeding on the host's blood.

Cats can become infected in several ways: by ingesting larvae from feces-contaminated material, by eating infected animals or insects, or through direct skin contact with larvae that penetrate the skin, especially around the paws. Unlike dogs, mother cats do not transmit hookworms through their milk.

Symptoms of hookworm infection vary depending on the severity. Adult cats may show few or no signs, but heavier infestations can cause diarrhea, weight loss, anemia (visible as pale gums), or black, tar-like stools due to digested blood. Other possible signs include weakness and poor body condition.

Hookworms pose a serious danger because they feed on blood. Kittens and young cats are especially vulnerable to severe illness and potentially fatal blood loss if left untreated.

Hookworms are also **zoonotic parasites**. Their larvae can penetrate human skin, causing intense itching and winding, red, tunnel-like marks known as *cutaneous larva migrans* or "ground itch." If swallowed, the larvae can also lead to intestinal problems in humans. Although hookworms are somewhat less common in cats than roundworms, their ability to cause anemia and skin

infections makes them an important parasite to prevent and control.

Tapeworms (Cestodes)

Tapeworms are segmented worms that live in the intestinal tract of cats. The most common species is *Dipylidium caninum*, also known as the cucumber seed tapeworm. The individual segments resemble grains of rice and can sometimes be seen in the cat's stool or around the anus.

Infection typically occurs when a cat ingests infected fleas during grooming or eats infected rodents. Fleas and rodents, in turn, become infected by swallowing tapeworm eggs from the environment.

Many cats with tapeworms show no obvious symptoms. Occasionally, mild digestive issues may appear, but the most visible sign is the presence of tapeworm segments near the cat's rear or in its litter box.

Tapeworm infections rarely cause serious illness in adult cats, but kittens may experience digestive upset or slowed growth. In severe cases, weight loss and nutrient deficiencies can occur.

Tapeworms are also **zoonotic**, though transmission to humans is rare. It usually happens when a person—often a child—accidentally ingests an infected flea. While the risk is low, this connection highlights the importance of consistent flea control as a key part of parasite prevention in cats.

Coccidia (Coccidia spp.)

Coccidia are microscopic, single-celled organisms (*Isospora spp*.) that cause coccidiosis. Almost all cats become exposed to *Isospora felis* at some point in their lives, usually as kittens.

Infection occurs when cats ingest oocysts (spore-like cysts) that are shed in feces and mature in the environment. Cats can also become infected by eating insects such as flies or cockroaches that carry the cysts.

In adult cats, coccidia infections rarely cause illness. However, in kittens, they can lead to serious disease because the parasites damage the intestinal lining, resulting in mucus-laden diarrhea, vomiting, and loss of appetite.

Coccidiosis can be dangerous for young cats, as severe infections may cause dehydration and even death. Fortunately, the *Isospora* species that infect cats generally do **not pose a risk to humans**, unlike *Toxoplasma gondii*. Coccidiosis mainly affects kittens, underlining how vulnerable young animals are to parasitic infections and how important hygiene and early treatment are for prevention.

Giardia (Giardia spp.)

Giardia are single-celled organisms that parasitize the small intestine of cats. The infection they cause, known as giardiasis, occurs in fewer than 5 percent of cats overall but can be significantly more common in multi-cat households, shelters, or breeding environments. Young cats under one year of age are especially susceptible.

Cats become infected by ingesting Giardia cysts from the feces of another infected animal—often a littermate—or from contaminated water sources.

Typical symptoms include foul-smelling diarrhea, vomiting, and weight loss. The diarrhea can be either acute or chronic, and some cats show no symptoms at all while still shedding infectious cysts.

Giardia infection can lead to dehydration and malnutrition, especially in kittens and cats with weakened immune systems. Certain genotypes of Giardia are **zoonotic** and can infect humans, causing gastrointestinal illness. Because of this, maintaining good hygiene and regularly cleaning litter boxes, food dishes, and living areas are important steps in preventing both feline and human infection.

Other Internal Parasites

In addition to the most common internal parasites, several other types can infect cats:

- **Stomach worms** (*Ollanulus tricuspis*, *Physaloptera spp.*): Infection occurs when a cat ingests vomit from an infected cat or eats intermediate hosts such as insects. Symptoms may include vomiting and loss of appetite.
- **Heartworms** (*Dirofilaria immitis*): Transmitted through mosquito bites. They can cause coughing, wheezing, breathing difficulties, vomiting, neurological problems, and, in severe cases, sudden death. Treatment in cats is difficult, making prevention essential.
- **Lungworms** (*Aelurostrongylus abstrusus*, *Eucoleus aerophilus*): Cats become infected by eating intermediate hosts (snails, slugs) or paratenic hosts (rodents, birds, lizards). Symptoms range from mild or no signs to coughing, wheezing, nasal discharge, and labored breathing.
- **Liver flukes:** Infection occurs through ingestion of infected intermediate hosts, such as land snails. These parasites typically affect the intestines, liver, bile ducts, and pancreatic ducts.
- **Bladder worms:** Occasionally infect cats and may cause urinary discomfort or blood in the urine.

These additional internal parasites require specific preventive and diagnostic measures, especially because many rely on intermediate hosts. The potentially severe consequences of infections such as heartworm and lungworm underline the importance of early detection, prevention, and veterinary guidance.

<u>Roundworms</u> (*Toxocara cati*)
Transmission: Ingestion of eggs from contaminated soil or water, consumption of infected rodents, or transmission through the mother's milk.
Symptoms: Often asymptomatic; may include diarrhea, vomiting, a swollen abdomen (especially in kittens), dull coat, weight loss, or coughing.
Risks: Malnutrition, digestive disorders, pneumonia caused by larval migration, anemia, and in severe cases, intestinal rupture.
Zoonosis: *Toxocara cati* can infect humans, causing visceral or ocular larva migrans, which may result in organ damage or blindness.

<u>Hookworms</u> (*Ancylostoma tubaeforme*)
Transmission: Ingestion of larvae from contaminated materials, infected animals, or insects; larvae can also penetrate the skin.
Symptoms: Often no visible signs; in heavier infestations, diarrhea, weight loss, anemia (pale gums), and dark, tar-like stools may occur.
Risks: Blood loss and severe anemia, especially dangerous for kittens.
Zoonosis: Larvae can penetrate human skin, causing *cutaneous larva migrans* ("ground itch") and sometimes intestinal discomfort.

<u>Tapeworms</u> (*Dipylidium caninum* and others)
Transmission: Ingestion of infected fleas or rodents.
Symptoms: Usually mild; visible rice-like segments near the anus or in feces, occasional digestive issues.

Risks: Digestive problems, delayed growth in kittens, and weight loss.
Zoonosis: Rare; most cases occur in children who accidentally swallow infected fleas.

Coccidia (*Isospora spp.*)
Transmission: Ingestion of oocysts from feces, contaminated soil, flies, or cockroaches.
Symptoms: Adult cats are usually symptom-free; kittens may experience mucous diarrhea, vomiting, loss of appetite, and dehydration.
Risks: Damage to the intestinal lining; severe infections in kittens can be fatal.
Zoonosis: Generally not transmissible to humans.

Giardia (*Giardia intestinalis*)
Transmission: Ingestion of cysts from feces or contaminated water.
Symptoms: Foul-smelling diarrhea, vomiting, and weight loss; some cats show no symptoms.
Risks: Dehydration and malnutrition, particularly in kittens or weakened cats.
Zoonosis: Certain strains can infect humans, causing giardiasis.

Heartworms (*Dirofilaria immitis*)
Transmission: Bite of infected mosquitoes.
Symptoms: Coughing, wheezing, shortness of breath, vomiting, neurological issues, and—in severe cases—sudden death.
Risks: Severe lung disease, heart failure, and organ damage.

Zoonosis: Not directly transmissible from cats to humans.

<u>Lungworms</u> (*Aelurostrongylus abstrusus*, *Capillaria aerophila*)
Transmission: Ingestion of intermediate hosts (snails, slugs) or paratenic hosts (rodents, birds, lizards).
Symptoms: Sometimes no signs; otherwise coughing, wheezing, nasal discharge, and breathing difficulties.
Risks: Bronchitis, pneumonia, and potentially fatal respiratory failure.

External Parasites
A Comprehensive Overview

External parasites live on the surface of a cat's body. They can cause irritation, skin problems, and transmit diseases. Even indoor cats are not completely safe from them. It's important to understand that external parasites are not just annoying—they can have serious health effects, and even cats that never go outdoors are at some level of risk.

Fleas (Siphonaptera)

Fleas are extremely common; most cats will experience at least one infestation in their lifetime. These small, jumping insects feed on the blood of their host. Infection occurs through contact with infested animals or contaminated environments. Fleas can also live on carpets, bedding, and furniture.

Typical symptoms of flea infestation include excessive scratching, licking, or biting of the skin, hair loss, flea dirt (tiny black specks resembling pepper), reddened or irritated skin, and tiny scabs known as miliary dermatitis.

For cats, fleas pose multiple health risks. Untreated infestations can cause severe itching, painful skin inflammation, and allergic reactions (flea allergy dermatitis). In heavy infestations—especially in kittens—fleas can cause anemia and even death. Fleas also transmit tapeworms (*Dipylidium caninum*) and other pathogens.

Humans can also be affected. Flea bites cause itching and skin irritation and can transmit diseases such as cat scratch disease (*Bartonella henselae*) and, rarely, plague. Because fleas reproduce quickly and can survive in the environment, effective control requires both treatment of the cat and thorough cleaning of the home.

Ticks (Ixodida)

Ticks are small, round arachnids that may appear black, pink, purple, or gray and are often mistaken for skin growths. These eight-legged parasites attach to the skin and feed on blood. They are more common in areas with abundant wildlife and among outdoor cats, being active year-round but especially during spring and fall.

Infection occurs when ticks attach to a cat while it's outside, particularly in grassy or wooded areas. Typical signs include visible ticks on the skin—most often around the head, neck, ears, and legs—and localized irritation or swelling at the bite site.

Ticks can transmit several serious diseases to cats, including Lyme disease, cytauxzoonosis ("bobcat fever," which can be fatal), mycoplasmosis, ehrlichiosis, and anaplasmosis. While cats are somewhat less likely than dogs to contract these diseases, they remain at risk.

Ticks also pose risks to humans, as they can spread Lyme disease, ehrlichiosis, Rocky Mountain spotted fever, and other infections. Because of these dangers, tick prevention is essential for outdoor cats. Regular tick checks and the proper removal technique—grasping the tick close to the skin with tweezers and pulling it out steadily—help prevent secondary infections.

Mites (Acari)

Mites are microscopic, spider-like parasites that can live on a cat's skin or in its ear canals.

Ear mites (*Otodectes cynotis*) are the most common and typically inhabit the ear canal, though they can spread to other areas of the body. They cause intense itching, leading to head shaking and scratching. Symptoms include dark, crumbly debris resembling coffee grounds, inflammation, and secondary ear infections. Kittens are particularly prone to ear mites.

Mange mites (*Notoedres cati*, *Sarcoptes scabiei*) cause mange or scabies. *Notoedres* infestations result in severe itching and inflamed skin, especially around the head and neck. *Demodex* mites live in hair follicles and sebaceous glands, causing patchy hair loss and mild irritation. Diagnosis is made through skin scrapings examined under a microscope.

Harvest mites (chiggers, *Trombiculidae*) are found in grassy or wooded areas and cause localized itching and irritation.

Mite infestations can lead to significant discomfort, secondary infections, and skin damage. Ear mites are highly contagious and can easily spread among cats in the same household. Some species, such as *Sarcoptes scabiei*, can cause temporary skin irritation in humans, though these infestations typically resolve on their own once the source is treated.

Lice (Phthiraptera)

Only one species of louse affects cats — *Felicola subrostratus*. Lice infestations are rare in well-cared-for cats and occur mostly in neglected, stray, or sick animals.

Transmission happens through direct contact with an infested cat. Symptoms include itching, visible lice or their eggs (nits) on the fur, and a rough or unkempt coat. In severe infestations, lice can cause skin irritation, restlessness, and even anemia due to blood loss.

Cat lice are **host-specific**, meaning they do not infest humans or other animals. Although they are much less common than fleas or mites, they can still cause discomfort and indicate that the affected cat may need medical care and improved grooming or nutrition.

General Signs of Parasitic Infestation What to Watch For

There are several general signs that may indicate a parasitic infection in cats:

- Unexplained weight loss
- Changes in appetite (increase or decrease)
- Vomiting
- Diarrhea (sometimes with mucus or blood)
- Frequent scratching, licking, or grooming
- Dull coat or hair loss
- Lethargy or decreased activity
- Visible parasites (worms or fleas) or their eggs in the stool or fur
- Swollen abdomen (especially in kittens with roundworms)
- Pale gums (a sign of anemia, especially with hookworms)
- Coughing or breathing difficulties (with lungworms or migrating larvae)
- Head shaking or ear scratching (with ear mites)

Because many of these symptoms can also occur with other health issues, a veterinary diagnosis is essential. Cat owners should pay attention to combinations of these signs and consult a veterinarian if any are observed.

Diagnosis by the Veterinarian
Methods and Procedures

Diagnosing a parasitic infection in cats usually begins with a thorough physical examination and a review of the cat's medical history. Depending on the suspected parasite, several diagnostic methods may be used:

- **Fecal examination:** A stool sample is analyzed under a microscope to detect parasite eggs (roundworms, hookworms, whipworms, tapeworms), cysts (Giardia, coccidia), or larvae (lungworms). Techniques such as direct smears, flotation (where eggs rise to the surface in a special solution), and sedimentation are commonly used. Since parasites are not shed continuously, multiple samples may be needed for accurate results.
- **Skin scraping:** Used to detect external parasites like mites (ear mites, mange mites, Demodex). A small amount of skin material is collected and examined microscopically.
- **Tapeworm detection:** Tapeworm segments are often visible to the naked eye around the anus or in feces. Flotation tests are not always reliable for tapeworm eggs.
- **Blood tests:** Primarily used to detect heartworm infections (antigen or antibody tests). Bloodwork can also reveal anemia or inflammation associated with parasites.

- **Additional tests:** In specific cases, other methods such as tracheal washes, bronchoalveolar lavage (for lungworms), or PCR testing (for Giardia or lungworms) may be performed.

Veterinarians have a range of tools for identifying parasites. Fecal tests are common for internal parasites, while skin scrapings help detect external ones. For certain infections, advanced diagnostics are essential to ensure accurate treatment and prevention.

Treatment Strategies
From Medication to Supportive Care

The treatment of parasitic infections in cats depends on the type of parasite involved and may include several approaches:

Medications:

- **Anthelmintics:** Drugs that kill internal worms (e.g., praziquantel for tapeworms, fenbendazole or pyrantel for roundworms and hookworms, and various medications for lungworms).
- **Insecticides/Acaricides:** Used to eliminate external parasites such as fleas, ticks, and mites (available as spot-on treatments, tablets, or collars containing ingredients like fipronil, imidacloprid, selamectin, or moxidectin).
- **Specific medications for protozoa:** For example, fenbendazole or metronidazole for giardiasis and certain antibiotics prescribed by veterinarians for coccidiosis.
- **Heartworm prevention:** Since treatment of heartworms in cats is limited, year-round preventive medication is strongly recommended.

Important: All medications should be prescribed by a veterinarian and administered exactly as directed. Some products formulated for dogs can be toxic to cats.

Supportive care:
In severe cases, hospitalization may be necessary for fluid

therapy (for dehydration), blood transfusions (for anemia), or oxygen therapy (for respiratory distress). Medications to control symptoms such as vomiting, diarrhea, or inflammation may also be prescribed.

Environmental control:
Thoroughly cleaning the home environment—especially sleeping areas and litter boxes—is essential to remove fleas, eggs, and larvae. Regular cleaning of litter boxes helps prevent the spread of eggs and cysts of internal parasites.

Treatment methods vary depending on the type of parasite. Veterinary guidance is crucial for choosing the right medication and dosage. In more severe cases, supportive therapy may be required to ensure full recovery.

Prevention Is the Best Protection
An Overview of Preventive Measures

Preventive care plays a vital role in protecting cats from parasitic infections:

Regular deworming:
Follow a deworming schedule recommended by your veterinarian, tailored to your cat's lifestyle and risk factors (for example, whether the cat goes outdoors or hunts).
Kittens require more frequent deworming, starting at a young age, and nursing queens should also be treated.

Flea and tick prevention:
Year-round use of veterinary-approved flea and tick preventives—such as spot-ons, oral medications, or collars—is highly effective. Always ensure the product is specifically designed for cats, as some dog products can be toxic.

Heartworm prevention:
Monthly administration of heartworm preventives should continue throughout the year, even for indoor cats, since mosquitoes can enter homes.

Hygiene measures:
Keep the litter box clean and remove feces daily to prevent the development of infectious parasite stages. Wash hands thoroughly after cleaning the litter box or handling the cat.

Discourage hunting and prevent cats from eating rodents or other wildlife. Provide well-cooked food and clean, fresh water.

Indoor vs. outdoor cats:
Indoor cats have a lower risk but are not immune to parasites. Outdoor cats face greater exposure through other animals and contaminated environments. Parasites can also be carried indoors by people or other pets.

Regular veterinary check-ups:
Annual or semi-annual examinations, including fecal tests, are essential for early detection and prevention. Discuss your cat's lifestyle with your veterinarian to develop an individualized parasite prevention plan.

Prevention is more effective—and often less expensive—than treatment. A combination of regular medication, good hygiene, and minimized risk factors is key. Even indoor cats benefit from consistent parasite protection.

Zoonotic Risks and Protective Measures for Cat Owners

Certain feline parasites can be transmitted to humans (zoonoses). Common zoonotic parasites include:

- **Roundworms (*Toxocara cati*):** Can cause visceral and ocular larva migrans in humans, leading to organ damage or blindness.
- **Hookworms (*Ancylostoma spp.*):** Larvae can penetrate human skin, causing *cutaneous larva migrans* ("ground itch").
- **Tapeworms (*Dipylidium caninum*):** Rarely transmitted, but possible through accidental ingestion of infected fleas, especially in children.
- **Giardia:** Some strains can infect humans, causing diarrhea and gastrointestinal discomfort.
- **Cat fleas (*Ctenocephalides felis*):** Can transmit *Bartonella henselae* (cat scratch disease).
- **Toxoplasma gondii:** A protozoan parasite that causes toxoplasmosis. It can be dangerous for pregnant women (leading to fetal illness or miscarriage) and immunocompromised individuals. Infection occurs through contact with contaminated feces or undercooked meat. Indoor cats pose a very low risk.
- **Lungworm (*Capillaria aerophila*):** Rarely, this parasite can cause respiratory illness in humans.

<u>To protect yourself from zoonotic infections</u>:

- Practice good hygiene — always wash your hands thoroughly with soap and water after touching your cat, before eating, and after cleaning the litter box.
- Wear gloves while gardening or handling soil, as it may contain parasite eggs.
- Prevent children from playing in areas where cats may defecate (like sandboxes), or ensure sandboxes are covered.
- Avoid direct contact with cat feces, especially during pregnancy or in cases of weakened immunity. If possible, have another person clean the litter box or wear gloves and wash hands immediately afterward.
- Control fleas both on the cat and in the home.
- Ensure your cat receives regular veterinary care, including parasite testing and preventive treatments.
- Cook meat thoroughly to kill *Toxoplasma* cysts.

Many cat parasites carry zoonotic potential, highlighting the importance of responsible pet ownership and hygiene to protect human health—especially for vulnerable groups.

Conclusion
A Healthy Life for Cats and Humans Through Parasite Control

In summary, parasites are common in cats and can cause a variety of health problems for both animals and humans. Proactive parasite control through regular veterinary care, preventive medication, proper hygiene, and environmental management is essential. With the right care, cats can live healthy, parasite-free lives, minimizing risks to their owners as well.

The responsibility for maintaining the health of both cats and humans lies equally with pet owners and veterinarians. By working together—through awareness, prevention, and timely treatment—we can ensure that our feline companions enjoy a safe, comfortable, and healthy life.

For your notes

For your notes